Journal of MOLECULAR BIOLOGY

EDITOR-IN-CHIEF
J. C. KENDREW

ASSISTANT EDITOR
S. BRENNER

EDITORIAL BOARD
M. S. Bretscher
C. R. Cantor
P. Chambon
W. Franke
M. Gellert
G. A. Gilbert
M. Gottesman
R. Huber
H. E. Huxley
A. Klug
V. Luzzati
M. F. Moody
K. Simons

VOLUME 148

5 May to 5 June 1981



ACADEMIC PRESS
London New York San Francisco

Copyright © 1981, by Academic Press Inc. (London) Ltd.

ALL RIGHTS RESERVED

No part of this volume may be reproduced in any form, by photostat, microfilm, or any other means, without written permission from the publishers.

Printed in Great Britain

CONTENTS OF VOLUME 148

NUMBER 1, 5 MAY 1981

GEISSELSODER, J., YOUDARIAN, P., DEHÒ, G., CHIDAMBARAM, M., GOLDSTEIN, R. and LJUNQUIST, E., Mutants of Satellite Virus P4 that Cannot Derepress their Bacteriophage P2 Helper 1

MENZEL, R. and ROTH, J., Regulation of the Genes for Proline Utilization in *Salmonella typhimurium*: Autogenous Repression by the *putA* Gene Product 21

SANCAR, A., WHARTON, R. P., SELTZER, S., KACINSKI, B. M., CLARKE, N. D. and RUPP, W. D., Identification of the *uvrA* Gene Product 45

SANCAR, A., CLARKE, N. D., GRISWOLD, J., KENNEDY, W. J. and RUPP, W. D., Identification of the *uvrB* Gene Product 63

BOGENHAGEN, D., LOWELL, C. and CLAYTON, D. A., Mechanism of Mitochondrial DNA Replication in Mouse L-cells. Replication of Unicircular Dimer Molecules 77

LETTERS TO THE EDITOR

RENKAWITZ-POHL, R., GLÄTZER, K. H. and KUNZ, W., Ribosomal RNA Genes with an Intervening Sequence are Clustered within the X Chromosomal Ribosomal DNA of *Drosophila hydei*. 95

MIZUNO, H., TOMITA, K.-I., NAKAGAWA, E., OHTSUKA, E. and IKEHARA, M., X-ray Diffraction Studies of the RNA Tetramer GpGpCpUp . . 103

NUMBER 2, 15 MAY 1981

BROSIUS, J., DULL, T. J., SLEETER, D. D. and NOLLER, H. F., Gene Organization and Primary Structure of a Ribosomal RNA Operon from *Escherickia coli* 107

EDEN, F. C., MUSTI, A. M. and SOBIESKI, D. A., Clusters of Repeated Sequences of Chicken DNA are Extensively Methylated but Contain Specific Undermethylated Regions 129

SILBERSTEIN, L. and LOWEY, S., Isolation and Distribution of Myosin Isoenzymes in Chicken Pectoralis Muscle 153

SCHUMAKER, V. N., POON, P. H., SEEGAN, G. W. and SMITH, C. A., Semi-flexible Joint in the C1q Subunit of the First Component of Human Complement 191

NUMBER 3, 25 MAY 1981

KRISCH, H. M. and SELZER, G. B., Construction and Properties of a Recombinant Plasmid Containing Gene *32* of Bacteriophage T4D . . 199

KARCH, F., TÖRÖK, I. and TISSIÈRES, A., Extensive Regions of Homology in Front of the Two hsp70 Heat Shock Variant Genes in *Drosophila melanogaster* 219

WILSON, M. C. and DARNELL, J. E., Jr, Control of Messenger RNA Concentration by Differential Cytoplasmic Half-life. Adenovirus Messenger RNAs from Transcription Units 1A and 1B 231

COHEN, F. E., STERNBERG, M. J. E. and TAYLOR, W. R., Analysis of the Tertiary Structure of Protein β-sheet Sandwiches 253

TREISMAN, R. and KAMEN, R., Structure of Polyoma Virus Late Nuclear RNA 273

NUMBER 4, 5 JUNE 1981

DUNN, J. J. and STUDIER, F. W., Nucleotide Sequence from the Genetic Left End of Bacteriophage T7 DNA to the Beginning of Gene *4* . . . 303

STILES, J. I., FRIEDMAN, L. R., HELMS, C., CONSAUL, S. and SHERMAN, F., Transposition of the Gene Cluster *CYC1-OSM1-RAD7* in Yeast . . 331

MYERS, R. M., WILLIAMS, R. C. and TIJAN, R., Oligomeric Structure of a Simian Virus 40 T Antigen in Free Form and Bound to DNA 347

DENIS, C. L., CIRIACY, M. and YOUNG, E. T., A Positive Regulatory Gene is Required for Accumulation of the Functional Messenger RNA for the Glucose-repressible Alcohol Dehydrogenase from *Saccharomyces cerevisiae* 355

BIEBRICHER, C. K., EIGEN, M. and LUCE, R., Product Analysis of RNA Generated *de novo* by Q_β Replicase 369

BIEBRICHER, C. K., EIGEN, M. and LUCE, R., Kinetic Analysis of Template-instructed and *de novo* RNA Synthesis by Q_β Replicase. 391

STEWART, M., Structure of α-tropomyosin Magnesium Paracrystals. II. Simulation of Staining Patterns from the Sequence and some Observations on the Mechanism of Positive Staining 411

LEDERER, F., GLATIGNY, A., BETHGE, P. H., BELLAMY, H. D. and MATHEWS, F. S., Improvement of the 2·5 Å Resolution Model of Cytochrome b_{562} by Redetermining the Primary Structure and Using Molecular Graphics . 427

COHEN, G. H., SILVERTON, E. W. and DAVIES, D. R., Refined Crystal Structure of γ-Chymotrypsin at 1·9 Å Resolution. Comparison with other Pancreatic Serine Proteases 449

LETTERS TO THE EDITOR

STAHL, S. J. and ZINN, K., Nucleotide Sequence of the Cloned Gene for Bacteriophage T7 RNA Polymerase 481

LANGE, R. H. and RICHTER, H.-P., A Symmetric Lipovitellin-Phosvitin Dimer in Cyclostome Yolk Platelet Crystals: Structural and Biochemical Observations 487

SIEKER, L. C., BOLLES, L., STENKAMP, R. E., JENSEN, L. H. and APPLEBY, C. A., Preliminary X-ray Study of a Dimeric Form of Hemerythrin from *Phascolosoma arcuatum* 493

Index to Volume 148

Adenovirus messenger RNAs from transcription units 1A and 1B; control of messenger RNA concentration by differential cytoplasmic half-life, 231
Alcohol dehydrogenase, glucose-repressible, from *Saccharomyces cerevisiae*, a positive regulatory gene is required for accumulation of the functional mRNA, 355
APPLEBY, C. A. (see also SIEKER, L. C.), 493

Bacteriophage P2 helper cannot be derepressed by mutants of satellite virus P4, 1
– P4 mutants that cannot derepress their bacteriophage P2 helper, 1
– T4D, construction and properties of a recombinant plasmid containing gene *32*, 199
— T7 DNA, nucleotide sequence from the genetic left end to the beginning of gene *4*, 303
— — RNA polymerase, nucleotide sequence of the cloned gene, 481
BELLAMY, H. D. (see also LEDERER, F.), 427
β-sheet sandwiches in proteins, analysis of the tertiary structure, 253
BETHGE, P. H. (see also LEDERER, F.), 427
BIEBRICHER, C. K., EIGEN, M. & LUCE, R., Kinetic analysis of template-instructed and *de novo* RNA synthesis by Q_β replicase, 391
– –, – – – –, Product analysis of RNA generated *de novo* by Q_β replicase, 369
BOGENHAGEN, D., LOWELL, C. & CLAYTON, D. A., Mechanism of mitochondrial DNA replication in mouse L-cells. Replication of unicircular dimer molecules, 77
BOLLES, L. (see also SIEKER, L. C.), 493
BROSIUS, J., DULL, T. J., SLEETER, D. D. & NOLLER, H. F., Gene organization and primary structure of a ribosomal RNA operon from *Escherichia coli*, 107

Chicken DNA, clusters of repeated sequences are extensively methylated but contain specific undermethylated regions, 129
— pectoralis muscle, isolation and distribution of myosin isoenzymes, 153
CHIDAMBARAM, M. (see also GEISSELSODER, J.), 1
γ-Chymotrypsin, refined crystal structure at 1·9 Å resolution; comparison with other pancreatic serine proteases, 449
CIRIACY, M. (see also DENIS, C. L.), 355
CLARKE, N. D. (see also SANCAR, A.), 45, 63
CLAYTON, D. A. (see also BOGENHAGEN, D.), 77
Cloned gene nucleotide sequence for bacteriophage T7 RNA polymerase, 481
Clusters of repeated sequences of chicken DNA are extensively methylated but contain specific undermethylated regions, 129
COHEN, F. E., STERNBERG, M. J. E. & TAYLOR, W. R., Analysis of the tertiary structure of protein β-sheet sandwiches, 253
—, G. H. SILVERTON, E. W. & DAVIES, D. R., Refined crystal structure of γ-chromotrypsin at 1·9 Å resolution. Comparison with other pancreatic serine proteases, 449
CONSAUL, S. (see also STILES, J. I.), 331
Construction and properties of a recombinant plasmid containing gene *32* of bacteriophage T4D, 199
C1q subunit of the first component of human complement has a semi-flexible joint, 191
CYC1-OSM1-RAD7 gene cluster, transposition in yeast, 331
Cyclostome yolk platelet crystals contain a symmetric lipovitellin-phosvitin dimer; structural and biochemical observations, 487
Cytochrome b_{562}, improvement of the 2·5 Å resolution model by redetermining the primary structure and using molecular graphs, 427
Cytoplasmic half-life, differential, and the control of mRNA concentration; adenovirus mRNAs from transcription units 1A and 1B, 231

DARNELL, J. E., JR (see also WILSON, M. C.), 231
DAVIES, D. R. (see also COHEN, G. H.), 449
DEHÒ, G. (see also GEISSELSODER, J.), 1
DENIS, C. L., CIRIACY, M. & YOUNG, E. T., A positive regulatory gene is required for accumulation of the functional messenger RNA for the glucose-repressible alcohol dehydrogenase from *Saccharomyces cerevisiae*, 355

DNA of bacteriophage T7, nucleotide sequence from the genetic left end to the beginning of gene *4*, 303
—, chicken, clusters of repeated sequences are extensively methylated but contain specific undermethylated region, 129
—, mitochondrial, mechanism of replication in mouse L-cells; replication of unicircular dimer molecules, 77
—, ribosomal, X chromosomal, of *Drosophila hydei* contains clustered ribosomal RNA genes with an intervening sequence, 95
Drosophila hydei, ribosomal RNA genes with an intervening sequence are clustered within the X chromosomal ribosomal DNA, 95
— *melanogaster*, extensive regions of homology in front of the two hsp70 heat shock variant genes, 219
DULL, T. J. (see also BROSIUS, J.), 107
DUNN, J. J. & STUDIER, F. W., Nucleotide sequence from the genetic left end of bacteriophage T7 DNA to the beginning of gene *4*, 303

EDEN, F. C., MUSTI, A. M. & SOBIESKI, D. A., Clusters of repeated sequences of chicken DNA are extensively methylated but contain specific undermethylated regions, 129
EIGEN, M. (see also BIEBRICHER, C. K.), 369, 391
Escherichia coli, gene organization and primary structure of a ribosomal RNA operon, 107

FRIEDMAN, L. R. (see also STILES, J. I.), 331

GEISSELSODER, J., YOUDARIAN, P., DEHÒ, G., CHIDAMBARAM, M., GOLDSTEIN, R. & LJUNQUIST, E., Mutants of satellite virus P4 that cannot derepress their bacteriophage P2 helper, 1
Gene, cloned, nucleotide sequence for bacteriophage T7 RNA polymerase, 481
— cluster *CYC1-OSM1-RAD7*, transposition in yeast, 303
— organization and primary structure of a ribosomal RNA operon from *Escherichia coli*, 170
—, positive regulatory, is required for accumulation of the functional mRNA for the glucose-repressible alcohol dehydrogenase from *Saccharomyces cerevisiae*, 355
— product, *uvrA*, identification, 45
— —, *uvrB*, identification, 63
— *32* of bacteriophage T4D contained in a recombinant plasmid; construction and properties of the plasmid, 199
Genes, hsp70 heat shock variant, in *Drosophila melanogaster*, are preceded by extensive regions of homology, 219
— for proline utilization, regulation in *Salmonella typhimurium*; autogeneous repression by the *putA* gene product, 21
—, ribosomal RNA, with an intervening sequence are clustered within the X chromosomal ribosomal DNA of *Drosophila hydei*, 95
GLATIGNY, A. (see also LEDERER, F.), 427
GLÄTZER, K. H. (see also RENKAWITZ-POHL, R.), 95
Glucose-repressible alcohol dehydrogenase from *Saccharomyces cerevisiae*, a positive regulatory gene is required for accumulation of the functional mRNA, 355
GOLDSTEIN, R. (see also GEISSELSODER, J.), 1
GpGpCpUp, X-ray diffraction studies of RNA tetramer, 103
GRISWOLD, J. (see also SANCAR, A.), 63

Half-life, cytoplasmic, differential, and the control of mRNA concentration; adenovirus mRNAs from transcription units 1A and 1B, 231
Heat shock variant genes, hsp70, in *Drosophila melanogaster*, are preceded by extensive regions of homology, 219
HELMS, C. (see also STILES, J. I.), 331
Helper, bacteriophage P2, cannot be depressed by mutants of satellite virus P4, 1
Hemerythrin from *Phascolosoma arcuatum*, preliminary X-ray study of a dimeric form, 493
Homologies, extensive, in front of the two hsp70 heat shock variant genes in *Drosophila melanogaster*, 219
hsp70 heat shock variant genes in *Drosophila melanogaster* are preceded by extensive regions of homology, 219
Human complement, semi-flexible joint in the C1q subunit of the first component, 191

IKEHARA, M. (see also MIZUNO, H.), 103

JENSEN, L. H. (see also SIEKER, L. C.), 493

Kacinski, B. M. (see also Sancar, A.), 45
Kamen, R. (see also Treisman, R.), 273
Karch, F., Török, I. & Tissières, A., Extensive regions of homology in fromt of the two hsp70 heat shock variant genes in *Drosophilia melanogaster*, 219
Kennedy, W. J. (see also Sancar, A.), 63
Kinetic analysis of template-instructed and *de novo* RNA synthesis by Q_β replicase, 391
Krisch, H. M. & Selzer, G. B., Construction and properties of a recombinant plasmid containing gene *32* of bacteriophage T4D, 199
Kunz, W. (see also Renkawitz-Pohl, R.), 95

Lange, R. H. & Richter, H.-P., A symmetric lipovitellin-phosvitin dimer in cyclostome yolk platelet crystals: structural and biochemical observations, 487
L-cells, mouse, mechanism of mitochondrial DNA replication; replication of unicircular dimer molecules, 77
Lederer, F., Glatigny, A., Bethge, P. H., Bellamy, H. D. & Mathews, F. S., Improvement of the 2·5 Å resolution model of cytochrome b_{562} by redetermining the primary structure and using molecular graphics, 427
Lipovitellin-phosvitin dimer, symmetrical, in cyclostome yolk platelet crystals: structural and biochemical observations, 487
Ljunquist, E. (see also Geisselsoder, J.), 1
Lowell, C. (see also Bogenhagen, D.), 77
Lowey, S. (see also Silberstein, L.), 153
Luce, R. (see also Biebricher, C. K.), 369, 391

Magnesium paracrystals of α-tropomyosin, structure; simulation of staining patterns from the sequence and some observations on the mechanism of positive staining, 411
Mathews, F. S. (see also Lederer, F.), 427
Menzel, R. & Roth, J., Regulation of the genes for proline utilization in *Salmonella typhimurium*: autogenous repression by the *putA* gene product, 21
Messenger RNA concentration control by differential cytoplasmic half-life; adenovirus messenger RNAs from transcription units 1A and 1B, 231
——, functional, for the glucose-repressible alcohol dehydrogenase from *Saccharomyces cerevisiae*, a positive regulatory gene is required for accumulation, 355
Methylation of clusters of repeated sequences of chicken DNA: the methylation is extensive but there are specific undermethylated regions, 129
Mitochondrial DNA replication mechanism in mouse L-cells; replication of unicircular dimer molecules, 77
Mizuno, H., Tomita, K.-I., Nakagawa, E., Ohtsuka, E. & Ikehara, M., X-ray diffraction studies of the RNA tetramer GpGpCpUp, 103
Mouse L-cells, mechanism of mitochondrial DNA replication; replication of unicircular dimer molecules, 77
Muscle, pectoralis, chicken, isolation and distribution of myosin isoenzymes, 153
Musti, A. M. (aee also Eden, F. C.), 129
Mutants of satellite virus P4 that cannot derepress their bacteriophage P2 helper, 1
Myers, R. M., Williams, R. C. & Tijan, R., Oligomeric structure of a simian virus 40 T antigen in free form and bound to DNA, 347
Myosin isoenzymes in chicken pectoralis muscle, isolation and distribution, 153

Nakagawa, E. (see also Mizuno, H.), 103
Noller, H. F. (see also Brosius, J.), 107

Ohtsuka, E. (see also Mizuno, H.), 103
Oligomeric structure of a simian virus 40 T antigen in free form and bound to DNA, 347

P2 (see also Bacteriophage), 1
P4 (see also Bacteriophage), 1
Phascolosoma arcuatum hemerythrin, preliminary X-ray study of a dimeric form, 493
Phosvitin-lipovitellin dimer, symmetric, in cyclostome yolk platelet crystals: structural and biochemical observations, 487
Plasmid containing gene *32* of bacteriophage T4D, construction and properties, 199
Polyoma virus late nuclear RNA, structure, 273
Poon, P. H. (see also Schumaker, V. N.), 191
Proline utilization in *Salmonella typhimurium*, regulation of the genes; autogenous repression by the *putA* gene product, 21

Protein β-sheet sandwiches, analysis of the tertiary structure, 253
putA gene product autogenous repression; regulation of the genes for proline utilization in *Salmonella typhimurium*, 21

Q_β replicase generation *de novo* of RNA; product analysis, 369
Q_β replicase synthesis of RNA, template-instructed and *de novo*; kinetic analysis, 391

Recombinant plasmid containing gene *32* of bacteriophage T4D, construction and properties, 199
Regulation of the genes for proline utilization in *Salmonella typhimurium*: autogenous repression by the *putA* gene product, 21
Regulatiory gene, positive, is required for accumulation of the functional mRNA for the glucose-repressible alcohol dehydrogenase from *Saccharomyces cerevisiae*, 355
Renkawitz-Pohl, R., Glätzer, K. H. & Kunz, W., Ribosomal RNA genes with an intervening sequence are clustered within the X chromosomal ribosomal DNA of *Drosophila hydei*, 95
Replication mechanism for mitochondrial DNA in mouse L-cells; replication of unicircular dimer molecules, 77
RNA generated *de novo* by Q_β replicase, product analysis, 369
—, late nuclear, polyoma virus-specific: structure, 273
— operon, ribosomal, from *Escherichia coli*, gene organization and primary structure, 107
— polymerase, bacteriophage T7, nucleotide sequence of the cloned gene, 481
—, ribosomal, genes with an intervening sequence are clustered within the X chromosomal ribosomal DNA of *Drosophila hydei*, 95
— synthesis by Q_β replicase, template-instructed and *de novo*; kinetic analysis, 391
— tetramer GpGpCpUp, X-ray diffraction studies, 103
Richter, H.-P. (see also Lange, R. H.), 487
Ribosomal RNA genes with an intervening sequence are clustered within the X chromosomal ribosomal DNA of *Drosophila hydei*, 95
— — operon from *Escherichia coli*, gene organization and primary structure, 107
Roth, J. (see also Menzel, R.), 21
Rupp, W. D. (see also Sancar, A.), 45, 63

Saccharomyces cerevisiae, a positive regulatory gene is required for accumulation of the functional mRNA for the glucose-repressible alcohol dehydrogenase, 355
Salmonella typhimurium, regulation of the genes for proline utilization; autogenous repression by the *putA* gene product, 21
Sancar, A., Clarke, N. D., Griswold, J., Kennedy, W. J. & Rupp, W. D., Identification of the *uvrB* gene product, 63
— —, Wharton, R. P., Seltzer, S., Kacinski, B. M., Clarke, N. D. & Rupp, W. D., Identification of the *uvrA* gene product, 45
Sandwiches, β-sheet, of proteins, analysis of the tertiary structure, 253
Satellite virus P4 mutants that cannot derepress their bacteriophage P2 helper, 1
Schumaker, V. N., Poon, P. H., Seegan, G. W. & Smith, C. A., Semi-flexible joint in the C1q subunit of the first component of human complement, 191
Seegan, G. W. (see also Schumaker, V. N.), 191
Seltzer, S. (see also Sancar, A.), 45
Selzer, G. B. (see also Krisch, H. M.), 199
Sequence, nucleotide, of the cloned gene for bacteriophage T7 RNA polymerase, 481
—, —, from the genetic left end of bacteriophage T7 DNA to the beginning of gene *4*, 303
Sequences, repeated, of chicken DNA, clusters are extensively methylated but contain specific undermethylated regions, 129
Serine proteases, pancreatic, comparison of structures; refined crystal structure of γ-chymotrypsin at 1·9 Å resolution, 449
Sherman, F. (see also Stiles, J. I.), 331
Sieker, L. C., Bolles, L., Stenkamp, R. E., Jensen, L. H. & Appleby, C. A., Preliminary X-ray study of a dimeric form of hemerythrin from *Phascolosoma arcuatum*, 493
Silberstein, L. & Lowey, S., Isolation and distribution of myosin isoenzymes in chicken pectoralis muscle, 153
Silverton, E. W. (see also Cohen, G. H.), 449
Simian virus 40 T antigen, oligomeric structure in free form and bound to DNA, 347
Sleeter, D. D. (see also Brosius, J.), 107
Smith, C. A. (see also Schumaker, V. N.), 191
Sobieski, D. A. (see also Eden, F. C.), 129

STAHL, S. J. & ZINN, K., Nucleotide sequence of the cloned gene for bacteriophage T7 RNA polymerase, 481
Staining pattern simulation and some observations on the mechanism of positive staining; structure of α-tropomyosin magnesium paracrystals, 411
STENKAMP, R. E. (see also SIEKER, L. C.), 493
STERNBERG, M. J. E. (see also COHEN, F. E.), 253
STEWART, M., Structure of α-tropomyosin magnesium paracrystals. II. Simulation of staining patterns from the sequence and some observations on the mechanism of positive staining, 411
STILES, J. I., FRIEDMAN, L. R., HELMS, C., CONSAUL, S. & SHERMAN, F., Transposition of the gene cluster *CYC1-OSM1-RAD7* in yeast, 331
Structural and biochemical observations of a symmetric lipovitellin-phosvitin dimer in cyclostome yolk platelet crystals, 487
Structure, crystal, of γ-chymotrypsin at 1·9 Å resolution; comparison with other pancreatic serine proteases, 449
— of cytochrome b_{562}, improvement of the 2·5 Å resolution model by redetermining the primary structure and using molecular graphs, 427
— — polyoma virus late nuclear RNA, 273
— — α-tropomyosin magnesium paracrystals; simulation of staining patterns from the sequence and some observations on the mechanism of positive staining, 411
—, oligomeric, of SV40 T antigen in free form and bound to DNA, 347
—, primary, and gene organization of a ribosomal RNA operon from *Escherichia coli*, 107
—, tertiary, of protein β-sheet sandwiches, an analysis, 253
STUDIER, F. W. (see also DUNN, J. J.), 303

T antigen, SV40, oligomeric structure in free form and bound to DNA, 347
T7 (see also Bacteriophage), 303, 481
T4D (see also Bacteriophage), 199
TAYLOR, W. R. (see also COHEN, F. E.), 253
TIJAN, R. (see also MYERS, R. M.), 347
TISSIÈRES, A. (see also KARCH, F.), 219
TOMITA, K.-I. (see also MIZUNO, H.), 103
TÖRÖK, I. (see also KARCH, F.), 219
Transcription units 1A and 1B, control of adenovirus mRNA concentration by differential cytoplasmic half-life, 231
— of the gene cluster *CYC1-OSM1-RAD7* in yeast, 331
TREISMAN, R. & KAMEN, R., Structure of polyoma virus late nuclear RNA, 273
α-Tropomyosin magnesium paracrystals, structure; simulation of staining patterns from the sequence and some observations on the mechanism of positive staining, 411

Unicircular dimer molecule replication; mechanism of mitochondrial DNA replication in mouse L-cells, 77
uvrA gene product identification, 45
uvrB gene product identification, 63

Virus P4 mutants that cannot derepress their bacteriophage P4 helper, 1
—, polyoma, structure of late nuclear RNA, 273
—, SV40, oligomeric structure of T antigen in free form and bound to DNA, 347

WHARTON, R. P. (see also SANCAR, A.), 45
WILLIAMS, R. C. (see also MYERS, R. M.), 347
WILSON, M. C. & DARNELL, J. E., JR, Control of messenger RNA concentration by differential cytoplasmic half-life. Adenovirus messenger RNAs from transcription units 1A and 1B, 231

X chromosomal ribosomal DNA of *Drosophila hydei* contains clustered ribosomal RNA genes with an intervening sequence, 95
X-ray diffraction studies of the RNA tetramer GpGpCpUp, 103
— study, preliminary, of a dimeric form of hemerythrin from *Phascolosoma arcuatum*, 493

Yeast, transposition of the gene cluster *CYC1-OSM1-RAD7*, 331
Yolk platelet crystals, cyclostome, contain a symmetric lipovitellin-phosvitin dimer; structural and biochemical observations, 487
YOUDARIAN, P. (see also GEISSELSODER, J.), 1
YOUNG, E. T. (see also DENIS, C. L.), 355

ZINN, K. (see also STAHL, S. J.), 481